

Elders Balmoral

2015 Drop Post Weaning and Yearling Assessment

Within-Site Results

Conducted by

Elders Balmoral Sire Evaluation Group



under the auspices of

The Australian Merino Sire Evaluation Association



Merino Lifetime Productivity Project Site



November 2016

Acknowledgement

The Merino Lifetime Productivity Project is being undertaken in partnership between the Australian Merino Sire Evaluation Association Incorporated (AMSEA) and Australian Wool Innovation (AWI). AMSEA and AWI would like to acknowledge those entities who also contribute funding, namely Woolgrowers through sire evaluation entry fees, site committee in-kind contributions, and sponsors of AMSEA. A special acknowledgement is also made to the Australian Government who support research, development and marketing of Australian wool.

Disclaimer

Australian Merino Sire Evaluation Association Incorporated (AMSEA) is funded by Australian Wool Innovation Limited (AWI) which gratefully acknowledges the funds provided by the Australian Government to support research, development and marketing of Australian wool. AMSEA sponsors, woolgrower entry fees and site committee in-kind contributions also contribute to AMSEA funding. The printing of this publication is funded by various advertisers and we thank them for their support. This publication should only be used as a general aid and is not a substitute for specific advice. To the extent permitted by law, AWI and AMSEA exclude all liability for loss or damage arising from the use of the information in this publication. © 2016 Australian Wool Innovation Limited and Australian Merino Sire Evaluation Association Incorporated. All rights reserved.

The Australian Merino Sire Evaluation Association has approved the format used in this report. Australian Sheep Breeding Values reported here are based on analyses conducted by Sheep Genetics.

Elders Balmoral Sire Evaluation Group Central Test Sire Evaluation

The Elders Balmoral Sire Evaluation Trials aim to evaluate and promote leading sires suited to fine wool production in Western Victoria.

This goal is achieved by informing participants, their clients and interested woolgrowers about the events surrounding the trials, and through producing and distributing annual reports and periodic newsletters. To further promote the evaluation, displays have been on show at the Australian Sheep & Wool Show now held in Bendigo, Balmoral Show and Hamilton Sheepvention.

Since 1999 successful annual open days have been held at “The Mountain Dam”, “Kerrsville”, “White Oaks”, “Arundale”, “Tuloona”, “Mokanger”, “Yiddinga”, “Wando Estate”, “Mepungah” and “Tuloona” to allow progeny inspections and to discuss the sire evaluation program with interested woolgrowers.

In 1998 a small group of stud breeders met to form what is now known as the Elders Balmoral Victoria Sire Evaluation Group. The Sire Evaluation Trials commenced in 1998 and as of this year there will be 19 progeny drops: 1998 - 2016. All trials are run for a minimum of 2 years. The site planning and direction is provided by the Elders Balmoral Sire Evaluation Management Committee.

Evaluations have been held on privately owned host properties around the Balmoral district progressing to a new property mostly every two years. Host properties run Merino fine wool ewes with genetics suitable for the district’s environment.

- 1998 & 1999 – “The Mountain Dam”, Balmoral
- 2000 & 2002 – “Kerrsville”, Balmoral
- 2002 & 2003 – “White Oaks”, Balmoral
- 2004 & 2005 – “Arundale”, Balmoral
- 2006 & 2007 – “Tuloona”, Harrow
- 2008 & 2009 – “Mokanger, Cavendish
- 2010 & 2011 – “Yiddinga”, Edenhope
- 2012 & 2013 – “Wando Estate”, Casterton
- 2014 – “Mepungah”, Wannon
- 2015 & 2016 – “Tuloona”, Harrow

Merino Lifetime Productivity Project

Over recent years we have used the base trials to value add with additional trials. An example of this is the fertility analysis of sires from the 2010 drop progeny, a pedigree collection comparison in 2012 and now the Merino Lifetime Productivity Project (MLP) trials in 2015 and 2016.

The MLP project is a partnership between AWI and AMSEA that aims to comprehensively explore lifetime relationships between wool production, carcase performance and fertility and compare early life measurements against lifetime performance. For modern Merino selection systems to be successful we need to understand and accommodate the lifetime relationships between all these production elements - and there are current data gaps. The MLP is set to fill these gaps through increasing our understanding of the genetics of the modern Merino sheep over its lifetime, across different locations and genotypes.

The Tuloona trial is one of five standard sire evaluation sites that will join via AI for two years and retain their ewe progeny for annual natural mating, classing and lifetime assessment. The sites will initially operate like a standard sire evaluation site – following the rigorous and independently measured and visual assessment protocols. At the conclusion of the standard sire evaluation (once progeny are 18-24 months of age) AWI will support the ongoing measurement and visual classing of ewe progeny through 4-5 joinings and annual shearings. The number of ewes AI’d to each sire is increased to 90 ewes to ensure that there will be sufficient ewe progeny numbers per sire throughout life. More MLP information is available at www.wool.com/MLP.

Thank you to our hosts, sponsors, committee and participants for enabling this valuable assessment of Merino genetics.

Tom Silcock

Chairman - Elders Balmoral Sire Evaluation Group

Contents

	Page
Foreword	1
Contents	2
Site Committee	3
Sire and Owner Details	4/5
Manager's Report	6
Assessment and Management Program	7
Visual Trait Assessment and Site Breeding Objective	8
Sire Codes and Pedigree	8
Summary Results	
Index Options	9
Table 1: AMSEA Index Values and Classer's Visual Grades	10
Figure 1a: Combined Measured and Visual Performance (DP+)	11
Figure 1b: Combined Measured and Visual Performance (MP+)	11
Figure 1c: Combined Measured and Visual Performance (FP+)	12
Figure 1d: Combined Measured and Visual Performance (WP+)	12
Figure 2: Fleece Weight and Fibre Diameter	13
Figure 3: Classer's Visual Grade: Tops and Culls	13
Figure 4: Fleece Weight and Body Weight	14
Figure 5: Fleece Weight and Fat	14
Figure 6: Fleece Weight and Eye Muscle Depth	15
Figure 7: Body Weight and Eye Muscle Depth	15
Figure 8: Staple Strength and Worm Egg Count	16
Detail Results	
Understanding the results – Measured trait performance	17
Table 2: Major Measured Traits and Classer's Visual Grade	18
Table 3: Other Measured Traits	19
Understanding the results – Visual trait performance	21
Table 4a: Wool Quality	21
Table 4b: Wool Quality and Pigmentation	22
Table 4c: Conformation	23
Table 4d: Breech	24
Other assessment results	
Table 5: Sire Means for Measured Traits	25
Understanding the results – Information to assist the use of results	
Accuracy of Flock Breeding Values (FBVs)	26
Link Sires	26
Calculation of combined information	26

2015 Drop Post Weaning and Yearling Assessment

The information in this Site Report provides an update of the assessment of the 2015 drop, including the Post Weaning and Yearling assessments of the sire's progeny performance for measured and visually assessed traits.

The Post Weaning fleece and visual assessments of the ewes were made at 7 months of age with 7 months of wool growth. Post Weaning shearing of the ewes was conducted at 7.5 months of age with 7.5 months of wool growth.

The Yearling fleece assessments and shearing of the wethers was conducted at 12 months of age with 12 months of wool growth. No visual assessment of the wethers was undertaken.

Updated Site Reports will be published annually, or when new information is available.

Site Committee

Tom Silcock (Chair)	03 5388 2238	tom@themountaindam.com.au
Mark Bunge (Deputy Chair)	03 5579 7224	bunge5@bigpond.com
Hugh Jarvis	03 5588 6356	hughjarvis@bigpond.com
Michael Craig (Treasurer and Host)	03 5588 1395	michael@tuloona.fastmail.com
Liz Mecham (Secretary)	0407 015 059	petelizmecham@bigpond.com
Nick Falkenberg	0407 559 680	nickfalk@optusnet.com.au
Tom Sweeny	0419 362 173	wandoestate@bigpond.com
Andrew Howells	0418 846 291	andrew.howells@elders.com.au
David Whyte	03 5572 2266	david.whyte@elders.com.au
Jim Farran	03 5585 1888	j.farran@bigpond.com
Richard Beggs	03 5577 8222	office@nareebnareeb.com.au
Richard McShane	03 5574 2367	mokanger2@bigpond.com
Tony Kealy	03 5586 5252	kealy6@bigpond.com
Steve Cotton	0447 352 321	s.cotton@dynamicag.com.au
Peter Mecham	0408 279 823	peter.mecham@zoetis.com
Sean Harvey	0417 869 674	sean@live.com.au
Dale Bruns	0458 899 918	dbruns@woolnetwork.com.au
Rosey Leeming	0408 704 243	meeblok1@bigpond.com
Russell Macgugan	0438 314 390	rmcgugan@woolnetwork.com.au
Scott Davis	0419 783 987	scott.davis@trutest.com.au
Hamish Dickson	0427 446 499	hamish@agripartner.com.au
Tom & Julia Waldron (Data collection)	03 5570 8359	minno.creek@gmail.com
Daniel Rogers (Data Collection)	03 5388 2257	danielr.yulong@gmail.com
Elise Kealy (Data Collection)	0448 600 525	elisekealy@gmail.com
Jonno Hicks (Data Collection)	03 5392 2366	jonno@hannaton.com.au
Anthony Close (Data Collection)	0413 285 860	anthonyclose92@gmail.com
Duncan Pixley (Data Collection)	0437 222 433	duncan.pixley@allflex.com.au
Amy Tierney (Data Collection)	0427 555 027	amy.tierney@elders.com.au

For further information on this report please contact

Tom Silcock:	03 5388 2238	tom@themountaindam.com.au
Ben Swain:	02 6743 2306	ben_swain@bigpond.com

Sire and Owner Details

Breeders flock, Sire name Sire ID #, Breed †	Owner Details
Billandri Poll, 130087 (Link) 600571-2013-130087, Poll Merino	Bill Sandilands Billandri, Kendenup WA 6323 P: (08) 9851 4030, F: (08) 9851 4264, E: csandilands@bournet.com.au
Bogo, 111424 (Link) 504792-2011-111424, Merino	Malcolm Peake Ravenswood, Boambolo Road, Yass NSW 2582 P: (02) 6227 1223, F: (02) 6227 1271, E: info@bogomerinos.com.au
Bundaleer Poll, 13V741 601449-2013-13V741, Poll Merino	Peter and Gavin Lieschke Pine Ridge, RMB 123, Walla Walla NSW 2659 P: (02) 6029 0142, F: (02) 6029 0188, E: lieschke@skymesh.com.au
Bundilla, 111265 504081-2011-111265, Merino	Ross, Rick & Jill Baldwin Bundilla, Tubbul Road, Young NSW 2594 P: (02) 6383 3802, F: (02) 6383 3805, E: bundillamerinos@bigpond.com
Centre Plus Poll, 207316 601250-2012-207316, Poll Merino	Robert Mortimer Devondale, Tullamore NSW 2874 P: (02) 6892 8259, F: (02) 6892 8292, E: robert@centreplus.com.au
Darriwell, 130941 503655-2013-130941, Merino	Russell Jones Darriwell, 924 Darriwell Rd, Trundle NSW 2875 P: (02) 6869 9242, F: (02) 6869 9242, E: darriwool@hotmail.com
Glenpaen, 120042 504654-2012-120042, Merino	Rod Miller 733 Victoria Valley Rd, Brimpaen VIC 3401 P: (03) 5383 9227, F: (03) 5383 9227, E: glenpaen@harboursat.com.au
Greenfields Poll, 130599 600240-2013-130599, Poll Merino	Daryl Smith Glenville, Cowell SA 5602 P: (08) 8628 5032, F: (08) 8628 5036, E: glenville@activ8.net.au
Hazeldean, 11.43 500383-2011-000043, Merino	Jim Litchfield Hazeldean Pty Ltd, Cooma NSW 2630 P: (02) 6453 5555, F: (02) 6453 5526, E: admin@hazeldean.com.au
Kurra-Wirra, SR5681 504173-2013-SR5681, Merino	Robert Close Kurra Wirra, 770 Moree-Culla Rd, Coleraine VIC 3315 P: (03) 5570 4238, F: (03) 5570 4234, E: kurrawirra@skymesh.com.au
Leahcim Poll, 090918 (Link) 600815-2009-090918, Poll Merino	Andrew and Rosemary Michael PO Box 31, Snowtown SA 5520 P: (08) 8865 2085, F: (08) 8865 2585, E: leahcimgenetics@bigpond.com
Leahcim Poll, 123153 600815-2012-123153, Poll Merino	Andrew and Rosemary Michael PO Box 31, Snowtown SA 5520 P: (08) 8865 2085, F: (08) 8865 2585, E: leahcimgenetics@bigpond.com
Merinotech WA Poll, 100081 (Unreg) 609040-2010-100081, Poll Merino	Ian Robertson Merinotech (WA) Ltd, RMB 311, Kojonup WA 6395 P: (08) 9833 6251, F: (08) 9833 6255, E: yarrak311@optusnet.com.au
Mokanger, 120092 (Link) 504888-2012-120092, Merino	Richard McShane Mokanger Past Co, 711 Mokanger Road, Cavendish VIC 3314 P: (03) 5574 2367, F: (03) 5574 2328, E: mokanger2@bigpond.com
Moojepin, 100248 (Link) 504637-2010-100248, Merino	Mark Wootton 1874 Hensley Park Road, Hensley Park VIC 3315 P: (03) 5574 8246, F: (03) 5574 8262, E: office@jigsawfarms.com.au

Sire and Owner Details

Breeders flock, Sire name Sire ID #, Breed [†]	Owner Details
Mumblebone, 130389 (Link) 500063-2013-130389, Merino	Chad Taylor Marapana, 456 Wuuluman Road, Wellington NSW 2820 P: (02) 6845 3620, F: (02) 6845 3608, E: chad@mumblebone.com.au
Mumblebone, 130850 500063-2013-130850, Merino	Chad Taylor Marapana, 456 Wuuluman Road, Wellington NSW 2820 P: (02) 6845 3620, F: (02) 6845 3608, E: chad@mumblebone.com.au
Nareeb Nareeb, 130380 500246-2013-130380, Merino	Richard Beggs 4395 Hamilton Chatsworth Rd, Nareeb VIC 3293 P: (03) 5577 8222, F: (03) 5577 8362, E: office@nareebnareeb.com.au
Nerstane, 130467 503298-2013-130467, Merino	John, Hamish and Jock McLaren Nerstane, Woolbrook NSW 2354 P: (02) 6777 5881, F: (02) 6777 5922, E: jock@nerstane.com.au
One Oak No. 2, R56 (Link) 503855-2010-100R56, Merino	Graham Wells 1763 Great Alpine Road, Smoko VIC 3741 E: oneoakpl@bigpond.com
Roseville Park, 140019 (Link) 504166-2014-140019, Merino	Matthew and Cherie Coddington Glenwood, 39R Dilladerry Rd MS3, Dubbo NSW 2830 P: (02) 6887 7286, E: rpmerinos@bigpond.com
The Mountain Dam, 11/ESA004 504572-2011-ESA004, Merino	Tom and Alison Silcock The Mountain Dam, 429 Silcocks Road, Telangatuk East VIC 3401 P: (03) 5388 2288, F: (03) 5388 2235, E: tom@themountaindam.com.au
Tuckwood Poll, 121021 601053-2012-121021, Poll Merino	Geoff Tucker PMB 21, Millicent SA 5280 P: (08) 8734 2050, F: (08) 8734 2052, E: geomag@activ8.net.au
Yalgoo, 120043 (Link) 501552-2012-120043, Merino	Jock Nivison Yalgoo, PO Box 141, Walcha NSW 2354 P: (02) 6777 2088, E: yalgoopartnership@bigpond.com
Yiddinga, 130374 (Unreg) 509242-2013-130374, Merino	Jim Farran PO Box 222, Edenhope VIC 3318 P: (03) 5585 1888, E: j.farran@bigpond.com

(Link) Sire evaluated to provide links between years and sites so that the all site results can be combined into a single report, e.g., *Merino Superior Sires*.

(Unreg) Sire bred in an unregistered flock.

- # Sire ID provides a unique number for all sheep. A sire ID has 16 digits.
- 2 for the breed of the flock, e.g., Merino (50), Poll Merino (60), Dohne (51), SAMM (48), Afrino (AF)
 - 4 for flock code, AASMB Registered flock code or unregistered code.
 - 4 for year of drop.
 - 6 for tag number used in the breeder's records.

[†] Breed of flock in which the sire was born

Host Property for 2015 drop progeny and location

“Tuloona” is operated by the Craig family and is located approximately 5km out of Harrow. Tuloona receives a winter dominated rainfall of approximately 470mm annually.

Ewe Base

The ewe base is a traditional super fine wool flock that has focussed over the past ten years on improving growth rate, wool cut and fertility whilst attempting to retain micron and quality. The mature ewe flock averages 17.2um and cuts 38kg/ha of clean wool (4.7kg at 63% yield, 2.96kg CFW) and weighing 52kg.

Ewes for the Merino Lifetime Productivity project were selected from three age groups totalling 3,500 ewes. Selection was based on evenness.

Joining

Laparoscopic insemination of 2160 ewes was conducted by Genstock Jerilderie between the 23-26 March, 2015. 25 sires were inseminated. Ewes averaged approximately condition score 2.9 at joining.

Owing to a mix up of semen at an AI centre, the semen sent for the Mumblebone sire was from two different Mumblebone sires. Both sires are reported in this report and the intended sire entrant has been re-joined and will have additional progeny in the 2016 drop.

Pregnancy and lambing

The ewes were pregnancy scanned on the 19 May, 2015. The results of which were disappointing with a relative low total number of foetuses. This was largely driven by a very low number of twin bearing ewes being scanned. In total, 67.3% foetuses to ewes joined were scanned.

Ewes were split following pregnancy scanning with the single bearing ewes divided into heavy condition and light condition in order to maximise lamb survival. It was also decided to lamb the single bearing ewes in groups of approximately 200 to increase the possibility of lamb survival.

Ewes completed lambing at Tuloona in late August 2015. A total of 1268 lambs were tagged on 3 September. This represented 87% of the number of foetuses scanned. DNA samples were taken at tagging to determine sire and dam parentage.

The lambs were marked at tagging and scored for breech traits. On 9 November the lambs were weaned with an average weaning weight of 23.4kg.

Weaning to Yearling Assessment

Lambs tracked well for growth rate and body weight gain from weaning until affected with lupinosis in March. Body weights collected early March averaged 31.2kg (up from 25.5 on 28/1/16 – 160 grams/day) with an average condition score of 2.98. A combination of managing the lupinosis and a number of trial activities further impacted on growth as outlined in the appended report by Hamish Dickson, AgriPartner Consulting. With shearing completed and green feed now in front of them, all lambs did well. With WEC, FAT & EMD all now collected the ewes are on track to meet their targeted joining weight.

Seasonal conditions

The Spring leading up to the Summer of 2016, gave us record rainfall deficiencies leading to many livestock water storages being dry for the first time ever. As a result, lambs were heavily supplemented from weaning to May. Record low levels of dry matter and water availability have now been replaced with an abundance of both. The wet season has now provided a big challenge to breakdown in wool and wool quality. An unscheduled fly protection application has been deemed imperative.

Assessment and Management Program

Activity		Date/s	Age	Wool
Selection of ewes		February 2015		
Allocation of ewes for mating		March 2015		
Pregnancy scanning		19 May 2015		
Allocated to lambing paddocks		10 August 2015		
Lambing: start – finish		16 – 23 August 2015		
Lambing mobs boxed to one management group		3 September 2015	14 days	
Tagging, pigmentation and breech scoring		3 September 2015	14 days	
Marking		3 September 2015	14 days	
Weaning		9 November 2015	81 days	
Mid side fleece sampling	P	17 March 2016	7 months	7 months
	Y (wethers)	17 August 2016	12 months	12 months
Visual trait scoring	P	17 March 2016	7 months	7 months
Shearing	P	11 April 2016	7.5 months	7.5 months
	Y (wethers)	17 August 2016	12 months	12 months
Fat and eye muscle scanning	Y	26 September 2016	13 months	6 months
Worm egg count sampling	H	10 October 2016	13.5 months	
Body weighing	W	9 November 2015	81 days	
	P	28 January 2016	5 months	
	P	24 March 2016	7 months	
	Y (wethers)	17 August 2016	12 months	
	Y	26 September 2016	13 months	
Drench	Drenched at weaning.			
	Ewes drenched in September 2016 when WEC samples were collected.			
Fly treatment	Treated with Klik® at marking. Progeny are not mulesed.			
	Ewes received a body and breech application of Klik in September 2016.			
Supplementary feeding	Silage, Barley and Lupins post weaning			
Field day or public display of 2015 drop progeny	Field Day & Progeny Display– April 2016			
	Static display at Balmoral Show – March 2016			
	Next Field Day 24 March 2017			

Visual Trait Assessment and Site Breeding Objective

Visual trait assessment

Classer's Grade: Mr David Whyte, Elders Limited

Trait Scores: Committee

Site Breeding Objective used to assess the Visual Classer's Grades

The Breeding Objective used by the classer/s when selecting the Classers Tops, Flock and Cull grades is described below. The Breeding Objective for both measured and visual assessed traits was developed by the site committee in consultation with the classer prior to the grading.

Breeding Objective

The goal is to select sheep that are productive and well grown, with sound conformation and carrying heavy fine wool fleeces of good character, colour and nourishment suitable for the western Victorian environment.

Sire Codes and Pedigrees

Sire code	Breeders flock, Sire number	Sheep Genetics ID	Sire of Sire
1	Billandri Poll, 130087	600571-2013-130087	509605-2009-090122
2	Bogo, 111424	504792-2011-111424	Unknown
3	Bundaleer Poll, 13V741	601449-2013-13V741	Unknown
4	Bundilla, 111265	504081-2011-111265	504081-2009-090044
5	Centre Plus Poll, 207316	601250-2012-207316	601250-2009-907538 (Centre Plus Poll, 907538)
6	Darriwell, 130941	503655-2013-130941	503655-2011-000952
7	Glenpaen, 120042	504654-2012-120042	503298-2009-090910 (Nerstane, 090910)
8	Greenfields Poll, 130599	600240-2013-130599	600240-2010-100087
9	Hazeldean, 11.43	500383-2011-000043	600553-2007-070002 (Coromandel Poll, ET2)
10	Kurra-Wirra, SR5681	504173-2013-SR5681	504173-2010-SR4862
11	Leahcim Poll, 090918	600815-2009-090918	600815-2007-070319
12	Leahcim Poll, 123153	600815-2012-123153	600815-2010-101009
13	Merinotech WA Poll, 100081	609040-2010-100081	609040-2008-088578
14	Mokanger, 120092	504888-2012-120092	504888-2009-000004
15	Moojepin, 100248	504637-2010-100248	504637-2008-081206
16	Mumblebone, 130389	500063-2013-130389	601365-2009-090399
17	Mumblebone, 130850	500063-2013-130850	500063-2010-100186
18	Nareeb Nareeb, 130380	500246-2013-130380	503855-2011-BL0115
19	Nerstane, 130467	503298-2013-130467	503298-2010-100919 (Nerstane, 100919)
20	One Oak No. 2, R56	503855-2010-100R56	503855-2008-080004
21	Roseville Park, 140019	504166-2014-140019	601050-2009-090853 (Stockman Poll, 090853)
22	The Mountain Dam, 11/ESA004	504572-2011-ESA004	600792-2009-090576 (Mernowie Poll, 090576)
23	Tuckwood Poll, 121021	601053-2012-121021	601082-2008-081375
24	Yalgoo, 120043	501552-2012-120043	503298-2008-080121 (Nerstane, 080121)
25	Yiddinga, 130374	509242-2013-130374	509242-2011-000076

Index Options

A breeding index combines multiple measured traits into a single value that reflects a certain emphasis on these traits. It is important that you use an index that best matches the breeding objective and production system of the flock you are selecting for.

It is recommended that the performance of individual measured and visually assessed traits is used in conjunction with an index as selection indexes assist in making balanced selection decisions.

Site Reports present 4 indexes, DP+; MP+; FP+ and WP+. The first 3 of these indexes are the same as MERINOSELECT indexes of that name but account for the fact that direct reproduction records have not been captured by AMSEA sire evaluation. The WP+ index is unique to AMSEA.

Provided is the percentage contribution that each trait makes to economic gain in a commercial flock that uses an index for sire selection. Additionally, included for each index are the likely within-flock responses from using an index for 10 years. These responses are based on a ram breeding flock with a standard breeding program, no introduction of outside genetics and uses 35% of their selection emphasis on traits that are not in the index (such as visually assessed performance).

Dual Purpose Plus (DP+)

Based on a meat focused production system where surplus progeny are sold as lambs and a portion of ewes are joined to terminal sires. Large increase in body weight and carcase traits. Moderate increase in fleece weight. Maintain fibre diameter and staple strength. Moderate increase in reproduction.

Merino Production Plus (MP+)

Based on a balanced wool and meat production system where surplus progeny are sold as hoggets. Balanced emphasis on increasing fleece weight and reduction in fibre diameter. Moderate increase in body weight, with little change in reproduction.

Fibre Production Plus (FP+)

Based on a wool production system where wethers are retained, operating in an environment where worms cause economic losses. Large reduction in fibre diameter. Moderate increase in staple strength. Small reduction in WEC (if measured in the breeding program). Small increase in fleece weight. Little change in body weight and reproduction.

Wool Production Plus (WP+)

Based on the MP+ production system with a greater emphasis on increasing fleece weight, while maintaining fibre diameter and a moderate emphasis on increasing body weight.




Table 1. AMSEA Index Values and Classer's Visual Grade

The index values reported are based on measured traits FBV performance with varying emphasis on fleece weight, fibre diameter, body weight, staple strength and worm egg count. See 'Index Options' (page 9) for more information on the indexes presented in the table below.

The highest performing sires for each trait (trait leaders) are highlighted by shading. Each sire is listed for Classer's Visual Grade and the same four indexes at all site evaluations.

Breeders flock, Sire name	Number of progeny	AMSEA Index Values				Classer's Visual Grade	
		Dual Purpose Plus	Merino Production Plus	Fibre Production Plus	Wool Production Plus	Tops % P	Culls % P
Billandri Poll, 130087	41	91	98	97	105	16	-3
Bogo, 111424	50	101	96	102	92	-12	3
Bundaleer Poll, 13V741	54	88	87	92	87	-7	-7
Bundilla, 111265	37	133	119	99	129	28	-8
Centre Plus Poll, 207316	44	101	102	110	96	-19	22
Darriwell, 130941	49	91	102	99	104	5	14
Glenpaen, 120042	49	85	110	111	102	-12	7
Greenfields Poll, 130599	48	100	109	105	109	-10	22
Hazeldean, 11.43	56	120	117	108	118	28	-13
Kurra-Wirra, SR5681	48	63	76	85	80	-8	28
Leahcim Poll, 090918	59	91	78	88	77	-9	1
Leahcim Poll, 123153	40	110	95	100	90	-4	-5
Merinotech WA Poll, 100081	54	110	107	113	100	-1	-9
Mokanger, 120092	37	97	99	99	101	-28	14
Moojepin, 100248	48	142	110	95	121	-17	15
Mumblebone, 130389	29	94	88	94	86	-12	-5
Mumblebone, 130850	27	117	104	101	102	43	-28
Nareeb Nareeb, 130380	45	80	105	104	107	23	-11
Nerstane, 130467	44	81	97	95	100	-8	3
One Oak No. 2, R56	57	84	85	86	89	-5	5
Roseville Park, 140019	33	90	90	92	91	-4	-10
The Mountain Dam, 11/ESA004	57	108	98	100	98	-2	-15
Tuckwood Poll, 121021	48	125	128	113	129	4	-7
Yalgoo, 120043	57	99	97	107	89	13	-11
Yiddinga, 130374	48	102	101	102	99	-1	-5
Average performance	46	100	100	100	100	27	28

Figure 1a. Combined measured traits (DP+ index) and combined visually assessed traits for the site objective.




Figure 1b. Combined measured traits (MP+ index) and combined visually assessed traits for the site objective.




Figure 1c. Combined measured traits (FP+ index) and combined visually assessed traits for the site objective.




Figure 1d. Combined measured traits (WP+ index) and combined visually assessed traits for the site objective.




Figure 2. Fleece weight by fibre diameter (FBVs)

The graph describes performance for fleece weight on the side axis and fibre diameter on the bottom axis. Sires that are above average for fleece weight and below average fibre diameter are located in the top left hand quarter.




Figure 3. Classer's Visual Grade - Tops by Cull

The graph describes performance for Classer's Visual Tops Grade on the side axis and Culls Grade on the bottom axis. Sires that have above average Tops and below average Culls are in the top left hand quarter.




Figure 4. Fleece weight by body weight (FBVs)

The graph describes performance for fleece weight on the side axis and body weight on the bottom axis. Sires that are above average for fleece weight and above average for body weight are located in the top right hand quarter.




Figure 5. Fleece weight by fat (FBVs)

The graph describes performance for fleece weight on the side axis and fat depth on the bottom axis. Sires that are above average for fleece weight and above average for fat are located in the top right hand quarter.




Figure 6. Fleece weight by eye muscle depth (FBVs)

The graph describes performance for fleece weight on the side axis and eye muscle depth on the bottom axis.

Sires that are above average for fleece weight and above average for eye muscle depth are located in the top right hand quarter.




Figure 7. Body weight by eye muscle depth (FBVs)

The graph describes performance for body weight on the side axis and eye muscle depth on the bottom axis. Sires that are above average for body weight and above average for eye muscle depth are located in the top right hand quarter.





Figure 8. Summary Graphs – SS and WEC

Figure 8. Staple strength by worm egg count (FBVs)

The graph describes performance for staple strength on the side axis and worm egg count on the bottom axis.

Sires that are above average for staple strength and above average for worm egg count are located in the top left hand quarter.



Understanding the Results

Measured trait performance and Classer's Visual Grade – Tables 2 and 3

Breeders flock, Sire number:	Identity of the breeder's flock and the sire's number or name.	
Number of progeny:	The number of progeny a sire had at the most recent measured analysis. Average number of progeny is included in Table 1.	
Flock Breeding Values:	<p>Flock Breeding Values (FBVs) are Estimated Breeding Values (EBVs) calculated by Sheep Genetics for the sires evaluated in this report. Only data from this site evaluation is used in the calculation of these FBVs. FBVs describe the relative breeding value (genetic performance) of the sires (in this case based on the performance of their progeny). A sire's progeny will express half of their sire's FBV. FBVs do not necessarily reflect the sire's observed performance, which is a combination of both genetic and environmental influences. FBVs are an estimate of the genetic component of the sheep's performance.</p> <p>The highest performing sires for each trait (trait leaders) are highlighted by shading. Curvature is the possible exception when for many breeders the optimum score is in the middle of the range therefore trait leaders have not been highlighted.</p>	
Traits: Abbreviation, trait and the (units reported)	<p>GFW: Greasy fleece weight (percentage).</p> <p>CFW: Clean fleece weight (percentage).</p> <p>FD: Average fibre diameter (micron).</p> <p>WT: Body weight (kilograms).</p> <p>FDCV: Fibre diameter coefficient of variation (percentage).</p> <p>SL: Staple length (mm) at the mid-side.</p> <p>SS: Staple strength (N/ktex) at the mid-side.</p> <p>EMD: Eye muscle depth (mm) at the 'C' site.</p> <p>FAT: Fat depth (mm) at the 'C' site.</p> <p>CURV: Fibre curvature (degrees).</p> <p>WEC: Worm egg count (% deviation in worm burden of sire's progeny).</p>	
Age at assessment:	<p>W = Weaning - 42 to 120 days (6 weeks to 4 months of age).</p> <p>E = Early Post Weaning - 120 to 210 days (4 to 7 months of age).</p> <p>P = Post Weaning - 210 to 300 days (7 to 10 months of age).</p> <p>Y = Yearling - 300 to 400 days (10 to 13 months of age).</p> <p>H = Hogget - 400 to 540 days (13 to 18 months of age).</p> <p>A = Adult - 540 days or older (18 months and older).</p>	
Classer's Visual Grade:	<p>A classer grades all progeny as either Tops, Flocks or Culls based on their visual assessment of all traits relative to the site's Breeding Objective. The percentage deviation from the average of Tops and Culls is presented in this report. Average percentage of Tops and Culls for the entire drop is included in Table 1.</p> <p>Page 8 provides more detail on Classer's Visual Grade and the site's Breeding Objective.</p>	

Table 2. Major Measured Traits and Classer's Visual Grade

Breeder's flock, Sire name	Number of Progeny	Post Weaning Flock Breeding Values (deviations)						Classer's Visual Grade ¹	
		GFW % Y [^]	CFW % Y	FD µm Y	WT kg P			Tops % P	Culls % P
Billandri Poll, 130087	41	9	9	0.7	-0.1	-0.6	-2.1	16	-3
Bogo, 111424	50	-8	-10	-1.2	-0.8	-1.5	-0.6	-12	3
Bundaleer Poll, 13V741	54	-7	-9	-0.2	-0.3	0.4	-0.6	-7	-7
Bundilla, 111265	37	12	15	0.7	2.7	4.7	7.5	28	-8
Centre Plus Poll, 207316	44	-4	-5	-1.2	-1.4	-2.0	-1.4	-19	22
Darriwell, 130941	49	3	4	0.2	1.0	0.7	-0.2	5	14
Glenpaen, 120042	49	1	-2	-1.4	-0.6	-1.3	-2.0	-12	7
Greenfields Poll, 130599	48	6	8	-0.6	-0.8	-1.4	-2.0	-10	22
Hazeldean, 11.43	56	5	5	-0.3	1.6	2.8	5.2	28	-13
Kurra-Wirra, SR5681	48	-2	-2	-0.2	-2.9	-5.6	-7.1	-8	28
Leahcim Poll, 090918	59	-12	-12	-0.4	-0.6	-1.6	-2.3	-9	1
Leahcim Poll, 123153	40	-9	-9	-0.3	0.2	0.1	0.2	-4	-5
Merinotech WA Poll, 100081	54	1	2	1.0	-2.2	-2.8	-2.5	-1	-9
Mokanger, 120092	37	3	5	-0.4	-0.6	-1.8	-2.5	-28	14
Moojepin, 100248	48	10	10	2.3	3.2	6.6	9.7	-17	15
Mumblebone, 130389	29	-7	-7	0.7	-1.0	-1.5	-2.4	-12	-5
Mumblebone, 130850	27	-3	-3	1.3	2.6	4.8	4.5	43	-28
Nareeb Nareeb, 130380	45	5	5	0.5	1.2	2.0	0.3	23	-11
Nerstane, 130467	44	5	2	-0.5	-0.8	-2.0	-1.4	-8	3
One Oak No. 2, R56	57	-3	-4	-1.0	-1.3	-2.3	-2.6	-5	5
Roseville Park, 140019	33	-5	-6	0.3	-0.2	0.2	-0.3	-4	-10
The Mountain Dam, 11/ESA004	57	-2	0	-0.2	-0.2	-0.1	-0.8	-2	-15
Tuckwood Poll, 121021	48	11	13	0.6	2.9	4.5	5.4	4	-7
Yalgoo, 120043	57	-7	-8	-1.5	-1.3	-2.1	-2.6	13	-11
Yiddinga, 130374	48	-4	-3	1.1	-0.4	0.0	0.8	-1	-5

[^] W = Weaning (42 to 120 days); P = Post Weaning (210 to 300 days); Y = Yearling (300 to 400 days); H = Hogget (400 to 540 days); A = Adult (540 days and older)

¹ Classer's Visual Grade is expressed as the percentage deviation of average Tops% and Culls%.

Table 3. Other Measured Traits

Breeder's flock, Sire name	Number of progeny	Flock Breeding Values (deviations)						
		FDCV	SL	SS	CURV	FAT	EMD	WEC
		% Y [^]	mm Y	N/ktex Y	deg/mm Y	mm Y	mm Y	% H
Billandri Poll, 130087	41	1.1	0.1	-1.1	-8.4	-0.5	-0.2	-32
Bogo, 111424	50	0.2	-9.0	-3.7	8.7	0.0	0.3	-10
Bundaleer Poll, 13V741	54	-0.2	-9.7	-3.1	5.3	-0.4	-0.3	-3
Bundilla, 111265	37	0.5	3.6	-5.2	-1.4	1.2	0.9	34
Centre Plus Poll, 207316	44	-0.5	1.6	-1.0	1.3	-0.3	0.0	-35
Darriwell, 130941	49	0.4	-4.0	0.7	-0.2	-0.7	-0.9	34
Glenpaen, 120042	49	1.2	-11.7	-0.2	10.3	-1.4	-2.1	68
Greenfields Poll, 130599	48	1.9	-0.6	-1.0	-0.5	-0.4	-0.3	54
Hazeldean, 11.43	56	0.5	4.7	-2.3	2.2	0.2	-0.1	11
Kurra-Wirra, SR5681	48	1.8	-8.0	-4.1	0.5	-0.5	-0.8	-9
Leahcim Poll, 090918	59	-0.3	-0.2	-1.5	2.7	0.3	0.9	-19
Leahcim Poll, 123153	40	-1.1	5.5	1.6	-3.6	1.2	1.3	-13
Merinotech WA Poll, 100081	54	-2.7	9.4	14.3	-3.0	1.1	1.1	-28
Mokanger, 120092	37	1.5	-8.3	-3.2	1.5	0.0	0.2	12
Moojepin, 100248	48	-1.2	24.9	1.9	-13.0	1.5	2.0	-38
Mumblebone, 130389	29	-1.4	3.6	4.9	-5.3	0.3	0.7	-22
Mumblebone, 130850	27	-2.6	6.7	7.5	-2.4	0.3	0.7	-24
Nareeb Nareeb, 130380	45	0.3	0.6	0.9	-4.0	-1.6	-2.3	-20
Nerstane, 130467	44	0.8	2.4	-5.1	-0.7	-0.9	-1.5	50
One Oak No. 2, R56	57	2.5	-13.6	-8.9	11.2	-0.6	-0.2	42
Roseville Park, 140019	33	-0.3	0.8	0.2	-1.3	-0.1	-0.4	14
The Mountain Dam, 11/ESA004	57	0.1	4.3	-1.2	-3.5	0.3	1.1	-7
Tuckwood Poll, 121021	48	0.1	7.9	3.7	-7.1	0.0	-0.3	63
Yalgoo, 120043	57	-0.9	-8.4	-1.6	11.2	0.3	0.5	-22
Yiddinga, 130374	48	-1.6	-2.8	7.3	-0.1	0.5	0.0	-21

[^] W = Weaning (42 to 120 days); P = Post Weaning (210 to 300 days); Y = Yearling (300 to 400 days); H = Hogget (400 to 540 days); A = Adult (540 days and older).

Understanding the results

Visual trait performance – Tables 4a, 4b, 4c, 4d

The following description of trait scores is a summary of the detailed word and diagrammatical description of these scores in Version 2 (2013) of the Visual Sheep Scores booklet that is available free from AWI or at www.merinosuperiorsires.com.au

A deviation from the average trait score for all progeny is reported as well as the percentage of the sire's progeny recorded for each trait.

■ Fleece rot:	The severity of fleece rot from 1 (no fleece rot), 2 and 3 (bands of bacterial staining but no crusting), and 4 and 5 (bands of crusty fleece rot).
■ Wool colour:	Greasy wool colour scored from 1 (whitest) to 5 (yellow).
■ Wool character:	Definition and variation of crimp between and along the staple scored from 1 (well defined and regular) to 5 (undefined and large variation).
■ Dust penetration:	Degree of dust penetration from 1 (only tip <6%) to 5 (71 to 100% of staple).
■ Staple weathering:	The deterioration due to light and water from 1 (least, <6% of staple) to 5 (most, 71 to 100%) reflect the depth and degree of deterioration.
■ Staple structure:	The size and diameter of each staple from 1 (<6mm) to 5 (>30 mm).
■ Fibre pigmentation:	The percentage of dark fibres on any part of the sheep from 1 (0 pigmented fibres at any site) to 5 (71 to 100% pigmented fibres at one or more sites). This trait does not include random spot or recessive black.
■ Non-fibre pigmentation:	The percentage of pigmentation on the areas not shorn from 1 (0 pigmentation at any site) to 5 (71 to 100% pigmented area on one or more bare skin sites, and/or 71 to 100% of the total hoof area).
■ Recessive black: (Black)	Recessive black (black) is identified by relatively symmetrical markings on both sides of the face. There are two scores 1 (no recessive markings) and 5 (recessive markings). This trait does not include random spot or fibre pigmentation.
■ Random spot: (Spot)	Random spot (spot) is identified by rounded wool or hair spot/s, not symmetrical. There are two scores 1 (no spot/s) and 5 (spot/s). If both sides of the face or body are spotted the sheep should be scored as a recessive black.
■ Face cover:	Wool cover on the face scored from 1 (open face) to 5 (fully covered face).
■ Feet/Legs:	Conformation of feet and legs scored from 1 (very straight) to 5 (very angulated).
■ Body wrinkle:	The degree of body wrinkle from 1 (no wrinkle) to 5 (extensive wrinkle).
■ Jaw:	The alignment of the lower jaw and its teeth relative to the top jaw from 1 (very well aligned) to 5 (heavily undershot or overshot).
■ Back/Shoulder:	Conformation of the back and shoulder from 1 (very square) to 5 (very dipped or high).
■ Breech cover	Size of natural bare area around the breech from 1 (large) to 5 (no bare).
■ Crutch cover	Size of natural bare area in the pubic and groin from 1 (large) to 5 (no bare).
■ Breech wrinkle	Degree of wrinkle at the tail set and hind legs from 1 (nil) to 5 (extensive).
■ Dag	Degree of dag adhering to the breech and legs from 1 (nil) to 5 (extensive).
■ Urine	Degree of urine stained wool in the breech area, including the hind legs from 1 (nil) to 5 (extensive).

Table 4a. Visual trait assessments – Wool Quality

Visually assessed traits reported were scored at their latest assessment with the exception of pigmentation which was scored at marking (Spot updated on an ongoing basis) and breech traits recorded at marking time (or later in unmulesed flocks with the exception of Dag and Urine). Traits are reported as a deviation (Dev) from the average trait score for all progeny. The percentage of a sire's progeny assessed for each score is also reported. No adjustments are made to the data to improve the accuracy of the results as is the case with sire means or breeding values. For the majority of breeder's objectives a negative deviation would be considered favourable and the larger the deviation the better.

Breeder's flock, Sire name	Wool Quality - Post Weaning																							
	Fleece Rot						Wool Colour						Wool Character						Dust Penetration					
	Dev	1	2	3	4	5	Dev	1	2	3	4	5	Dev	1	2	3	4	5	Dev	1	2	3	4	5
Billandri Poll, 130087	0.0	94	6	0	0	0	0.2	0	13	75	12	0	0.1	0	19	75	6	0	-0.4	0	50	44	6	0
Bogo, 111424	0.0	100	0	0	0	0	0.2	4	15	58	23	0	-0.2	0	42	58	0	0	0.0	0	23	65	12	0
Bundaleer Poll, 13V741	0.0	100	0	0	0	0	-0.1	3	21	69	7	0	-0.2	0	41	55	4	0	-0.1	0	31	55	14	0
Bundilla, 111265	0.0	100	0	0	0	0	-0.1	0	45	40	15	0	0.2	0	10	80	10	0	0.1	0	15	70	15	0
Centre Plus Poll, 207316	0.0	100	0	0	0	0	0.2	0	16	67	17	0	-0.2	0	42	54	4	0	0.0	0	21	62	17	0
Darriwell, 130941	0.0	100	0	0	0	0	0.0	0	26	58	16	0	0.2	0	5	84	11	0	0.1	0	16	63	21	0
Glenpaen, 120042	0.0	100	0	0	0	0	0.3	0	8	69	23	0	-0.3	0	50	50	0	0	0.2	0	12	69	19	0
Greenfields Poll, 130599	0.0	95	5	0	0	0	0.0	4	14	77	5	0	0.0	0	27	68	5	0	0.2	0	18	50	32	0
Hazeldean, 11.43	0.0	100	0	0	0	0	0.0	0	33	48	19	0	-0.3	0	56	41	3	0	0.0	0	26	52	22	0
Kurra-Wirra, SR5681	0.1	92	8	0	0	0	0.2	0	16	60	24	0	-0.4	0	64	36	0	0	-0.1	0	36	44	20	0
Leahcim Poll, 090918	0.0	96	4	0	0	0	0.0	0	29	61	10	0	0.0	0	29	61	10	0	0.1	0	21	54	25	0
Leahcim Poll, 123153	0.0	100	0	0	0	0	0.0	0	23	68	9	0	-0.1	0	32	64	4	0	0.0	0	32	41	27	0
Merinotech WA Poll, 100081	0.0	100	0	0	0	0	-0.3	0	46	50	4	0	0.1	0	23	65	12	0	-0.3	0	50	42	8	0
Mokanger, 120092	0.0	95	5	0	0	0	0.1	0	20	65	15	0	-0.1	0	25	75	0	0	0.2	0	15	55	30	0
Moojepin, 100248	0.0	100	0	0	0	0	0.4	0	4	67	29	0	0.8	0	0	43	57	0	0.5	0	10	43	43	4
Mumblebone, 130389	0.0	100	0	0	0	0	-0.3	0	54	38	8	0	0.0	0	15	85	0	0	0.0	0	15	77	8	0
Mumblebone, 130850	0.1	92	8	0	0	0	0.2	0	31	38	31	0	0.1	0	8	92	0	0	0.1	0	23	54	23	0
Nareeb Nareeb, 130380	0.0	100	0	0	0	0	0.0	0	25	62	13	0	0.1	0	12	88	0	0	-0.4	0	46	54	0	0
Nerstane, 130467	0.0	96	4	0	0	0	-0.2	0	50	35	15	0	-0.2	0	35	65	0	0	0.2	0	19	50	31	0
One Oak No. 2, R56	0.0	100	0	0	0	0	0.1	0	22	61	17	0	-0.1	0	33	67	0	0	-0.1	0	31	56	13	0
Roseville Park, 140019	0.0	100	0	0	0	0	0.2	0	24	47	29	0	-0.2	0	35	65	0	0	0.0	0	18	76	6	0
The Mountain Dam, 11/ESA004	0.0	100	0	0	0	0	-0.1	0	35	55	10	0	0.2	0	10	77	13	0	-0.2	0	35	61	4	0
Tuckwood Poll, 121021	0.0	100	0	0	0	0	-0.3	0	52	41	7	0	0.3	0	3	83	14	0	0.1	0	17	62	21	0
Yalgoo, 120043	0.0	100	0	0	0	0	-0.5	0	63	37	0	0	-0.1	0	33	63	4	0	-0.2	0	33	63	4	0
Yiddinga, 130374	0.0	100	0	0	0	0	-0.2	0	37	59	4	0	0.2	0	11	74	15	0	-0.1	0	41	41	18	0
Average performance	1.0	98	2	0	0	0	2.8	0	29	56	15	0	2.8	0	26	67	7	0	2.9	0	26	56	18	0

Table 4b. Visual trait assessments – Wool Quality and Pigmentation

For the majority of breeder's objectives a negative deviation for wool quality traits would be considered favourable and the larger the deviation the better. Staple Structure is the possible exception when for many breeders the optimum score is in the middle of the range therefore trait leaders have not been highlighted. Four pigmentation traits are reported. Fibre pigmentation and Non-fibre pigmentation are scored **1** to **5**, however Recessive black and Random spot are scored **1** (no pigmentation of this type) or **5** (when the trait is expressed). Only the percentage progeny for each sire that a score 5 is recorded, are reported for Recessive black and Random spot.

Breeder's flock, Sire name	Wool Quality - Post Weaning												Pigmentation - Marking													
	Staple Weathering						Staple Structure						Fibre pigmentation						Non-fibre pigmentation					Black	Spot	
	Dev	1	2	3	4	5	Dev	1	2	3	4	5	Dev	1	2	3	4	5	Dev	1	2	3	4	5	5	5
Billandri Poll, 130087							0.1	7	12	81	0	0	0.0	98	2	0	0	0	0.1	33	61	6	0	0	0	2
Bogo, 111424							-0.3	0	58	42	0	0	0.0	94	4	0	2	0	0.0	43	48	9	0	0	0	0
Bundaleer Poll, 13V741							0.0	0	34	66	0	0	0.0	97	1	2	0	0	-0.2	59	41	0	0	0	0	0
Bundilla, 111265							0.2	0	15	85	0	0	0.0	97	0	0	0	3	0.3	21	67	12	0	0	0	5
Centre Plus Poll, 207316							-0.2	0	54	46	0	0	-0.1	100	0	0	0	0	-0.1	43	54	3	0	0	0	0
Darriwell, 130941							0.3	0	5	95	0	0	-0.1	100	0	0	0	0	0.3	26	53	21	0	0	0	0
Glenpaen, 120042							-0.4	0	69	31	0	0	-0.1	100	0	0	0	0	-0.3	69	30	1	0	0	0	2
Greenfields Poll, 130599							0.1	0	23	77	0	0	0.0	96	2	2	0	0	-0.3	65	35	0	0	0	0	0
Hazeldean, 11.43							-0.4	0	74	26	0	0	0.0	97	3	0	0	0	-0.1	48	48	4	0	0	0	0
Kurra-Wirra, SR5681							-0.2	0	56	44	0	0	0.0	95	4	1	0	0	-0.1	51	40	9	0	0	0	4
Leahcim Poll, 090918							0.1	0	25	75	0	0	0.0	97	1	2	0	0	-0.2	62	35	1	2	0	0	0
Leahcim Poll, 123153							0.0	0	27	73	0	0	0.0	93	5	2	0	0	0.0	45	45	10	0	0	0	0
Merinotech WA Poll, 100081							0.0	0	27	73	0	0	-0.1	100	0	0	0	0	-0.3	65	35	0	0	0	0	2
Mokanger, 120092							0.0	0	35	65	0	0	0.3	79	16	2	0	3	0.4	21	58	18	3	0	0	0
Moojepin, 100248							0.3	0	0	100	0	0	0.1	90	8	0	2	0	-0.1	55	39	6	0	0	0	4
Mumblebone, 130389							-0.1	0	38	62	0	0	0.1	91	6	0	0	3	0.3	28	50	19	3	0	0	0
Mumblebone, 130850							0.0	0	31	69	0	0	0.0	96	4	0	0	0	0.1	42	46	12	0	0	0	0
Nareeb Nareeb, 130380							0.1	0	17	83	0	0	0.0	96	2	2	0	0	0.3	25	56	19	0	0	0	2
Nerstane, 130467							-0.2	0	54	46	0	0	0.0	94	6	0	0	0	0.3	17	68	15	0	0	0	0
One Oak No. 2, R56							0.0	0	28	72	0	0	0.0	98	2	0	0	0	-0.3	63	37	0	0	0	0	2
Roseville Park, 140019							0.0	0	29	71	0	0	-0.1	100	0	0	0	0	0.2	23	69	8	0	0	0	0
The Mountain Dam, 11/ESA004							0.1	4	19	77	0	0	-0.1	100	0	0	0	0	0.0	44	47	7	2	0	0	0
Tuckwood Poll, 121021							0.2	0	10	90	0	0	-0.1	100	0	0	0	0	-0.3	61	39	0	0	0	0	0
Yalgoo, 120043							0.0	0	27	73	0	0	-0.1	100	0	0	0	0	-0.4	78	22	0	0	0	0	0
Yiddinga, 130374							0.3	0	4	96	0	0	0.0	96	4	0	0	0	0.3	10	84	6	0	0	0	0
Average performance							2.7	0	31	69	0	0	1.1	96	3	1	0	0	1.6	44	48	8	0	0		

Table 4c. Visual trait assessments – Conformation

Traits are reported as a deviation (Dev) from the average trait score for all progeny. The percentage of a sire's progeny assessed for each score is also reported. No adjustments are made to the data to improve the accuracy of the results as is the case with sire means or breeding values.

For the majority of breeder's objectives a negative deviation would be considered favourable and the larger the deviation the better. Face cover is the possible exception when for many breeders the optimum score is in the middle of the range therefore trait leaders have not been highlighted.

Breeder's flock, Sire name	Conformation - Post Weaning																													
	Jaw						Legs and Feet						Shoulder and Back						Face Cover						Body Wrinkle					
	Dev	1	2	3	4	5	Dev	1	2	3	4	5	Dev	1	2	3	4	5	Dev	1	2	3	4	5	Dev	1	2	3	4	5
Billandri Poll, 130087	0.0	100	0	0	0	0	0.1	24	38	38	0	0	0.1	62	38	0	0	0	-0.1	0	25	75	0	0	0.2	0	50	50	0	0
Bogo, 111424	0.1	96	0	4	0	0	0.1	19	54	27	0	0	0.0	75	25	0	0	0	0.3	0	0	85	15	0	0.1	0	62	38	0	0
Bundaleer Poll, 13V741	0.0	100	0	0	0	0	-0.1	24	62	14	0	0	-0.1	83	17	0	0	0	0.0	0	14	83	3	0	0.0	3	59	38	0	0
Bundilla, 111265	0.0	100	0	0	0	0	-0.2	40	40	20	0	0	-0.1	84	16	0	0	0	0.2	0	0	90	10	0	0.2	0	47	53	0	0
Centre Plus Poll, 207316	0.0	100	0	0	0	0	0.2	13	58	29	0	0	-0.1	83	17	0	0	0	-0.3	0	46	54	0	0	0.1	4	48	48	0	0
Darriwell, 130941	0.0	100	0	0	0	0	-0.3	37	58	5	0	0	0.2	63	32	0	5	0	0.3	0	5	74	21	0	0.2	0	63	26	11	0
Glenpaen, 120042	0.0	100	0	0	0	0	0.0	23	58	19	0	0	0.3	58	31	11	0	0	0.1	0	15	73	12	0	0.3	0	46	42	12	0
Greenfields Poll, 130599	0.0	100	0	0	0	0	0.5	5	45	50	0	0	0.1	68	23	9	0	0	0.1	0	5	95	0	0	0.1	5	50	45	0	0
Hazeldean, 11.43	0.0	100	0	0	0	0	-0.2	30	63	7	0	0	0.1	63	37	0	0	0	-0.1	0	30	67	3	0	0.2	0	44	56	0	0
Kurra-Wirra, SR5681	0.1	96	0	4	0	0	0.5	12	28	60	0	0	0.4	48	39	13	0	0	0.1	0	12	76	12	0	0.2	0	57	39	4	0
Leahcim Poll, 090918	0.0	100	0	0	0	0	-0.1	43	28	29	0	0	0.1	70	26	4	0	0	-0.1	0	25	71	4	0	-0.6	26	74	0	0	0
Leahcim Poll, 123153	0.0	100	0	0	0	0	0.1	9	73	18	0	0	-0.1	82	18	0	0	0	-0.2	0	36	64	0	0	-0.5	27	68	5	0	0
Merinotech WA Poll, 100081	0.0	100	0	0	0	0	-0.1	15	77	8	0	0	0.0	73	27	0	0	0	-0.2	0	35	65	0	0	0.2	0	46	54	0	0
Mokanger, 120092	0.0	100	0	0	0	0	0.3	10	55	35	0	0	0.1	68	26	6	0	0	0.2	0	0	90	10	0	0.5	0	37	47	16	0
Moojepin, 100248	0.0	100	0	0	0	0	-0.1	29	57	14	0	0	-0.2	90	10	0	0	0	-0.6	0	71	29	0	0	-0.6	29	71	0	0	0
Mumblebone, 130389	0.0	100	0	0	0	0	-0.1	38	31	31	0	0	-0.2	92	8	0	0	0	0.1	0	0	100	0	0	-0.2	0	85	15	0	0
Mumblebone, 130850	0.0	100	0	0	0	0	-0.3	38	54	8	0	0	0.0	77	15	8	0	0	-0.5	0	62	38	0	0	-0.8	46	54	0	0	0
Nareeb Nareeb, 130380	0.0	100	0	0	0	0	-0.2	38	50	12	0	0	-0.2	88	12	0	0	0	-0.1	4	21	75	0	0	0.2	0	50	46	4	0
Nerstane, 130467	0.1	96	0	4	0	0	0.3	7	58	35	0	0	0.0	69	31	0	0	0	0.3	0	0	81	19	0	0.3	0	35	65	0	0
One Oak No. 2, R56	0.0	100	0	0	0	0	-0.2	33	53	14	0	0	0.1	75	11	14	0	0	0.2	0	3	86	11	0	0.3	0	44	50	6	0
Roseville Park, 140019	0.0	100	0	0	0	0	-0.1	24	65	11	0	0	0.0	82	12	6	0	0	-0.1	0	24	76	0	0	0.0	6	53	41	0	0
The Mountain Dam, 11/ESA004	0.0	100	0	0	0	0	0.1	26	39	35	0	0	0.1	77	6	13	4	0	0.1	0	6	94	0	0	-0.2	4	77	19	0	0
Tuckwood Poll, 121021	0.0	100	0	0	0	0	0.2	21	41	38	0	0	-0.1	86	14	0	0	0	-0.1	0	28	72	0	0	-0.3	11	79	10	0	0
Yalgoo, 120043	0.0	100	0	0	0	0	-0.4	43	50	7	0	0	-0.1	80	20	0	0	0	0.1	0	10	83	7	0	0.3	0	40	60	0	0
Yiddinga, 130374	0.0	100	0	0	0	0	0.1	22	52	26	0	0	-0.2	96	4	0	0	0	0.0	0	15	85	0	0	-0.2	15	59	26	0	0
Average performance	1.0	100	0	0	0	0	2.0	25	51	24	0	0	1.3	76	21	3	0	0	2.9	0	19	75	6	0	2.3	7	56	35	2	0

Table 4d. Visual trait assessments – Breech

Traits are reported as a deviation (Dev) from the average trait score for all progeny. The percentage of a sire's progeny assessed for each score is also reported. No adjustments are made to the data to improve the accuracy of the results as is the case with sire means or breeding values.

For the majority of breeder's objectives a negative deviation would be considered favourable and the larger the deviation the better.

Breeders flock, Sire name	Breech Visual Traits																													
	Breech Cover						Breech Wrinkle						Breech Cover						Breech Wrinkle						Dag					
	Marking						Marking						Post Weaning						Post Weaning						Post Weaning					
	Dev	1	2	3	4	5	Dev	1	2	3	4	5	Dev	1	2	3	4	5	Dev	1	2	3	4	5	Dev	1	2	3	4	5
Billandri Poll, 130087	0.0	0	12	60	28	0	0.2	6	42	40	12	0	0.0	0	17	65	18	0	0.1	0	53	41	6	0	-0.3	81	19	0	0	0
Bogo, 111424	-0.1	0	19	57	22	2	0.0	11	43	41	5	0	-0.1	0	21	62	17	0	-0.1	0	67	33	0	0	-0.1	64	28	8	0	0
Bundaleer Poll, 13V741	-0.3	7	24	52	17	0	-0.2	16	53	24	5	2	0.2	0	18	48	34	0	0.1	4	48	48	0	0	0.3	45	31	21	3	0
Bundilla, 111265	0.0	3	15	55	25	2	0.2	5	42	38	12	3	-0.1	0	25	60	15	0	0.1	5	50	40	5	0	-0.3	75	25	0	0	0
Centre Plus Poll, 207316	-0.3	5	30	41	24	0	-0.2	11	59	24	6	0	-0.2	0	35	52	13	0	0.3	5	39	43	13	0	-0.2	70	26	4	0	0
Darriwell, 130941	0.1	0	8	62	26	4	0.2	11	30	43	13	3	0.2	0	21	37	42	0	0.3	0	42	47	11	0	-0.3	79	16	5	0	0
Glenpaen, 120042	0.2	0	9	54	31	6	0.6	1	28	46	19	6	0.2	0	23	31	46	0	0.4	0	35	46	19	0	0.0	65	27	4	0	4
Greenfields Poll, 130599	0.0	0	14	59	25	2	0.0	16	41	34	7	2	-0.1	8	9	61	22	0	0.1	5	43	48	4	0	0.4	41	41	0	18	0
Hazeldean, 11.43	0.1	0	19	45	32	4	0.4	1	31	55	13	0	0.1	0	15	63	22	0	0.2	3	41	52	4	0	-0.1	69	23	8	0	0
Kurra-Wirra, SR5681	0.1	0	14	49	33	4	0.5	2	28	42	28	0	0.1	0	18	52	30	0	0.2	0	48	48	4	0	-0.4	88	12	0	0	0
Leahcim Poll, 090918	-0.4	3	34	48	15	0	-0.8	46	46	8	0	0	-0.2	4	33	44	19	0	-0.6	22	74	4	0	0	-0.3	86	11	3	0	0
Leahcim Poll, 123153	0.0	4	16	45	30	5	-0.5	30	50	20	0	0	-0.1	0	30	52	18	0	-0.6	30	61	9	0	0	0.3	45	32	23	0	0
Merinotech WA Poll, 100081	-0.2	5	26	46	20	3	-0.1	15	44	33	8	0	0.1	0	11	62	27	0	0.1	0	46	54	0	0	0.3	38	42	15	5	0
Mokanger, 120092	0.3	0	2	50	45	3	0.5	3	29	39	29	0	0.1	0	16	58	26	0	0.4	0	26	63	11	0	0.7	30	30	30	10	0
Moojepin, 100248	-0.4	4	30	50	16	0	-0.6	32	52	16	0	0	-0.4	0	43	52	5	0	-0.5	14	86	0	0	0	-0.4	86	14	0	0	0
Mumblebone, 130389	0.2	0	9	53	34	4	-0.1	16	47	31	6	0	-0.1	0	31	46	23	0	0.0	0	62	38	0	0	-0.2	69	31	0	0	0
Mumblebone, 130850	-0.3	0	33	52	15	0	-0.9	52	44	4	0	0	-0.5	7	40	53	0	0	-1.1	73	27	0	0	0	-0.3	80	20	0	0	0
Nareeb Nareeb, 130380	0.0	0	16	51	31	2	-0.2	18	41	37	4	0	0.0	0	28	44	28	0	0.3	0	36	60	4	0	0.0	60	32	8	0	0
Nerstane, 130467	0.5	0	8	32	47	13	0.6	3	25	40	28	4	0.6	0	0	46	50	4	0.5	0	19	73	8	0	0.4	42	35	15	8	0
One Oak No. 2, R56	0.7	1	3	29	48	19	1.0	0	14	41	37	8	0.1	0	16	56	28	0	0.3	0	39	53	8	0	-0.1	72	19	6	3	0
Roseville Park, 140019	-0.1	3	11	64	22	0	-0.4	31	42	25	2	0	0.0	0	16	67	17	0	0.2	6	39	44	11	0	-0.1	72	22	0	6	0
The Mountain Dam, 11/ESA004	0.1	2	16	44	36	2	-0.2	21	43	33	3	0	0.0	0	22	50	28	0	-0.3	9	72	19	0	0	-0.1	65	32	0	3	0
Tuckwood Poll, 121021	-0.1	2	31	37	22	8	-0.1	20	37	35	8	0	0.2	0	14	55	28	3	-0.3	0	86	14	0	0	0.1	59	26	7	4	4
Yalgoo, 120043	0.1	2	10	56	27	5	0.3	10	24	46	20	0	0.1	0	13	58	29	0	0.2	0	45	48	7	0	0.2	50	33	10	7	0
Yiddinga, 130374	-0.1	0	19	54	27	0	-0.3	17	52	31	0	0	-0.1	0	22	59	19	0	-0.2	14	56	30	0	0	0.5	37	33	22	4	4
Average performance	3.1	2	17	50	28	3	2.4	16	39	33	11	1	3.0	2	21	53	24	0	2.4	8	50	38	4	0	1.5	63	26	8	3	0

Table 5. Sire Means for Measured Traits

Sire means are the average performance of all the progeny of a sire adjusted for all available information on sex, birth type, rear type, age of dam, age of measurement and management group, in order to improve the accuracy. No account is made for trait heritability and genetic correlations between traits that can improve the breeding value accuracy, as is the case in Table 1.

The highest performing sires for each trait (trait leaders) are highlighted by shading. Curvature is the possible exception when for many breeders the optimum score is in the middle of the range therefore trait leaders have not been highlighted. The **Progeny group average** listed at the bottom of the table is the actual mean of the progeny group.

Breeder's flock, Sire name	Number of Progeny	Sire means for measured traits (deviations from the site mean)										
		GFW	CFW	FD	FDCV	SL	SS	WT			FAT	EMD
		kg Y [^]	kg Y	µm Y	% Y	mm Y	N/ktex Y	W	kg P	Y	mm Y	mm Y
Billandri Poll, 130087	41	0.1	0.0	0.2	1.0	-2.7	-1.7	0.1	0.0	-1.3	-0.1	0.1
Bogo, 111424	50	-0.1	-0.1	-0.5	0.1	-6.9	-3.8	-0.3	-1.4	-0.1	0.0	0.1
Bundaleer Poll, 13V741	54	-0.1	-0.2	0.1	-0.4	-8.8	-5.1	-0.5	0.3	-0.4	-0.1	-0.3
Bundilla, 111265	37	0.3	0.2	0.3	0.1	1.5	-6.2	1.2	1.7	4.2	0.3	0.3
Centre Plus Poll, 207316	44	0.0	-0.1	-0.7	-0.2	2.1	-1.2	-0.8	-1.6	-0.6	-0.1	0.1
Darriwell, 130941	49	0.1	0.1	0.1	0.2	-3.8	0.4	0.9	0.5	-0.1	-0.2	-0.7
Glenpaen, 120042	49	0.1	0.0	-0.7	0.8	-8.8	1.1	-0.2	-0.6	-0.8	-0.2	-1.1
Greenfields Poll, 130599	48	0.2	0.2	-0.4	1.4	0.1	1.7	-0.5	-0.4	-1.1	-0.1	-0.1
Hazeldean, 11.43	56	0.0	0.1	0.0	0.2	5.0	-0.6	0.7	0.4	3.1	0.0	-0.3
Kurra-Wirra, SR5681	48	-0.1	0.0	0.0	1.1	-7.2	-4.0	-1.1	-3.1	-3.3	-0.1	-0.5
Leahcim Poll, 090918	59	-0.2	-0.1	-0.3	0.0	0.8	-1.9	0.1	-0.4	-1.0	0.0	0.6
Leahcim Poll, 123153	40	-0.2	-0.1	-0.5	-0.1	4.7	1.3	0.2	0.6	-0.2	0.3	0.7
Merinotech WA Poll, 100081	54	0.1	0.1	0.6	-1.5	8.6	14.8	-1.4	-1.1	-1.4	0.3	0.8
Mokanger, 120092	37	0.2	0.2	0.0	0.6	-7.6	-1.6	0.1	-0.8	-1.4	0.0	0.4
Moojepin, 100248	48	0.2	0.1	1.2	-0.7	20.5	1.8	0.8	3.2	5.6	0.3	1.0
Mumblebone, 130389	29	-0.2	-0.1	0.4	-0.9	2.8	4.8	-0.7	-0.1	-2.1	0.1	0.5
Mumblebone, 130850	27	-0.1	-0.1	0.6	-1.8	6.7	6.2	1.3	3.4	1.2	-0.1	0.3
Nareeb Nareeb, 130380	45	0.0	0.0	0.3	0.5	-0.1	-0.8	0.7	1.2	-0.5	-0.3	-1.4
Nerstane, 130467	44	0.1	0.0	0.0	0.1	3.2	-5.5	-0.1	-1.9	0.5	-0.2	-0.9
One Oak No. 2, R56	57	0.0	0.0	-0.5	1.6	-11.7	-9.1	-0.6	-1.3	-0.9	-0.1	0.0
Roseville Park, 140019	33	-0.2	-0.2	0.1	-0.5	0.8	0.0	-0.4	0.4	-0.3	0.0	-0.3
The Mountain Dam, 11/ESA00	57	0.0	0.0	-0.2	-0.2	4.1	0.2	-0.1	0.5	-1.0	0.0	0.6
Tuckwood Poll, 121021	48	0.2	0.2	0.3	0.2	6.7	6.0	1.7	2.2	2.8	0.0	-0.3
Yalgoo, 120043	57	-0.1	-0.1	-0.9	-0.5	-6.5	-3.3	-0.3	-1.1	-1.5	0.1	0.4
Yiddinga, 130374	48	-0.1	0.0	0.7	-1.0	-3.6	6.5	-0.6	-0.7	0.6	0.1	-0.1
Progeny group average	46	2.6	1.7	14.3	20.1	78.9	25.8	23.4	28.4	32.6	2.6	21.1
		kg	kg	µm	%	mm	N/ktex	kg			mm	mm

Accuracy of Flock Breeding Values

Flock Breeding Values (FBVs) are reported by Sheep Genetics (SG). FBVs express the expected performance of progeny of a sire relative to another sire in the evaluation when mated to the same standard of ewes. FBVs improve the accuracy of sire results because they account for the association between traits, adjustment for birth effects and the number of progeny a sire has in the analysis.

True Breeding Values would be achieved if the number of progeny evaluated for each sire were infinite. Because the number of progeny in the evaluation is not infinite, performance shown in this report is described as *Flock* Breeding Values.

Without progeny test information the correlation between the *Flock* and *True* Breeding Value of sires from different sources would be zero (0.0%). The correlation between *Flock* and *True* Breeding Value improves rapidly from 0.0% with no progeny to 77% with 10 progeny. The rate of improvement in correlation slows from 86% with 20 progeny, to 90% with 30 progeny and 92% with 40 progeny. With an infinite population the correlation is 100%. Note that the correlation used in the above example is for a trait such as fibre diameter with a high heritability (0.5).

A heritability of 0.5 indicates that half or 50% of the measured performance is passed onto offspring. A heritability of 0.35 indicates 35% is passed on. The FBVs that are shown in this report have already accounted for heritability and therefore describe the performance that can be expected from a sire's progeny.

Link Sires

Link sires provide the 'genetic link' between sire evaluation sites located across Australia to allow all sires entered in these site evaluations to have their performance reported relative to each other in Merino Superior Sires. Merino Superior Sires reports sires from across all effectively linked sire evaluation sites and across all evaluations at these sites. Link sires are therefore a vital component of the sire evaluation.

To be used as a link a sire must have at least 25 progeny assessed at 1st Assessment at one accredited site. Site reports provide valuable information not reported in Merino Superior Sires however Merino Superior Sires reports the performance of a large number of sires which can provide a wider perspective of the elite sires available across many flocks in Australia.

Calculation of Combined Information

Combined measured trait performance is calculated as Index – 100. Three different index options are provided to cater for breeders' different breeding objectives.

Combined visual trait performance is calculated as:

$(\text{Classer's Visual Grade Tops\%} - \text{Culls\%})/5$, expressed as a deviation from $(\text{average Tops\%} - \text{average Culls\%})/5$.

Example

Sire's performance: \square AMSEA DP+ Index value = 119.7
 \square Tops% = 25.5 (average Tops% = 25.1)
 \square Culls% = 17.6 (average Culls% = 16.4)

Combined Measured = $119.7 - 100 = 19.7$
Combined Visual = $((25.5 - 17.6)/5) - ((25.1 - 16.4)/5)$
 = $7.9/5 - 8.7/5 = 1.58 - 1.74 = -0.1$

ELDERS BALMORAL SIRE EVALUATION GROUP

ANNUAL FIELD DAY

FRIDAY, MARCH 24, 2017

“Tuloona”

4580 Coleraine-Edenhope Road HARROW VIC

Sheep classing workshop

How to best use the tools of the trade

Visual assessments ASBVs Ramselect



45 progeny groups on display

2015 and 2016 Merino Lifetime

Productivity project sire progeny

Autodrafting demonstration

Industry displays and demonstrations

Lunch and refreshments provided

www.balmoralbreeders.com.au

www.wool.com/MLP

For more information:

Tom Silcock — 0419 882 238 tom@themountainindam.com.au; Liz Mecham — 0407 015 059 petelizmecham@bigpond.com

**Balmoral
Breeders**



Elders Balmoral

2015 Drop

Post Weaning and Yearling Assessment



Merino Lifetime Productivity Project Site

